Surgery performed on:

**RIGHT EYE** 

LEFT EYE

## **IOL Post-Operative Drop Instructions**

Please bring <u>all</u> post-operative drops to your 1 day and 1 week appointment for review.

3 in 1 Combination Drops – to be provided by your Patient Counselor  Drop name – Prednisolone/Gatifloxacin/Bromfenac Ophthalmic Solution  Cap color – Yellow			
Instruc Week #	tions: Start date: End 1: apply 1 drop 3 times per day in the eye tha	date: at had surgery.	
Week #2: apply 1 drop 2 times per day in the eye that had surgery.			Prednisolone P.O.,   Catiflexacin  fromfenac Ophthalmic Solution 1% / 0.5% / 0.075%
$\square$ Week #3: apply 1 drop 2 times per day in the eye that had surgery. Then discontinue.			O WILL
Week #4: after 21 days drop should be discontinued. See instruction above.			
OR	ual Duana - ta ha fillad and manidad huusaw	u Dhawaa ay	
	ual Drops – to be filled and provided by your		
**You must pick up <b>all 3 drops</b> from your pharmacy prior to cataract surgery**  **Apply drops a minimum of 3 minutes apart**			
	Drop name – Ofloxacin or Besivance Cap color – Tan		Offloxacin Ophthalmic Solution, USP 0.3% Servin Ophthalmic
	Instructions: Start date:		Solution  5 mL Sterile Sterile Contractorial
2.	Drop Name – Lotemax Gel or Prednisolone A Cap color – Pink or white	Acetate or Durezol	NDC 61314-637-15
	Instructions: Start date:		Prednisolone Acetate Ophthalmic Suspension USP
	Week #2 – use 1 drop 2 times per day in the	eye that had surgery. SHAKE BOTTLE.	1% R only
	Week #3 – use 1 drop 2 times per day in the	eye that had surgery. SHAKE BOTTLE.	NOC NUMBERON AND THE PROPERTY AND A SECTION AND A SECTION ASSESSMENT AND A SECTION ASSESSMENT ASSES
	Then discontinue.		The LOTEMAX's proposed on the control of the contro
3.	. Drop Name – Diclofenac or Ketorolac or Prolensa Cap color – Grey		
	Instructions: Start date:		VOC SCHOOL WID CO.
	Week #1 – use 1 drop 3 times per day in the	, ,	BAUSCH+LONB
	Week #2 – use 1 drop 3 times per day in the	PROLENSA' perofine contraine	
	Week #3 – use 1 drop 3 times per day in the eye that had surgery.		solution/007%
	Then discontinue.  **If you have Prolensa, use drop 1 time per day in the eye that had surgery for 4 weeks.		

<sup>\*</sup>If you have new pain or a sudden decrease in vision please call DLV or your co-managing optometrist  $^{\ast}$