

Surgery performed on:
RIGHT EYE LEFT EYE

ICL Post-Operative Drop Instructions

Please bring all post-operative drops to your 1 day and 1 week appointment for review.

2 in 1 Combination Drops – to be provided by your Patient Counselor

1. Drop name – Prednisolone/Gatifloxacin Ophthalmic Solution
Cap color – Grey

Instructions: Start date: _____ End date: _____
*Week #1 – apply 1 drop 3 times per day in the eye(s) that had surgery.
Then discontinue.*



2. Drop Name – Alphagan P or Brimonidine
Cap color – Green

Instructions: Start date: _____ End date: _____
*Week #1 – apply 1 drop 3 times per day in the eye(s) that had surgery.
Then discontinue.*



--OR--

Individual Drops – to be filled and provided by your Pharmacy

****You must pick up all 3 drops from your pharmacy prior to ICL surgery****

****Apply drops a minimum of 3 minutes apart****

1. Drop name – Ofloxacin or Besivance
Cap color – Tan

Instructions: Start date: _____ End date: _____
*Week #1 – use 1 drop 3 times per day for 1 week in the surgical eye(s).
Then discontinue.*



2. Drop Name – Lotemax Gel or Prednisolone Acetate
Cap color – Pink or white

Instructions: Start date: _____ End date: _____
*Week #1 – apply 1 drop 3 times per day in the eye(s) that had surgery.
SHAKE BOTTLE. Then discontinue.*



3. Drop Name – Alphagan P or Brimonidine
Cap color – Green

Instructions: Start date: _____ End date: _____
*Week #1 – apply 1 drop 3 times per day in the eye(s) that had surgery.
Then discontinue.*



If you have new pain or a sudden decrease in vision please call DLV or your co-managing optometrist